

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE (DD-MM-YYYY)		2. REPORT TYPE Conference Proceeding		3. DATES COVERED (From - To) 18-22 November 2002	
4. TITLE AND SUBTITLE Variability in the Backscattering to Scattering and F/Q Ratios Observed in Natural Waters				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER 601153N	
				5d. PROJECT NUMBER	
6. AUTHOR(S) Sherwin D. Ladner, Robert A. Arnone, Richard W. Gould, Alan D. Weidemann, Vladimir I. Haltrin, A. P. Lee, Paul M. Martinolich				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Research Laboratory Oceanography Division Stennis Space Center, MS 39529-5004				8. PERFORMING ORGANIZATION REPORT NUMBER NRL/PP/7330/02/0062	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Naval Research 800 N. Quincy St. Arlington, VA 22217-5660				10. SPONSOR/MONITOR'S ACRONYM(S) ONR	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution is unlimited					
20030523 107					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Variability of the backscattering to scattering (bb/b) and the upwelling irradiance/upwelling radiance (f/Q) ratios affect the accurate retrieval of inherent optical properties from ocean color satellite algorithms (SeaWiFS and MODIS). We investigate the variability of bb/b and the f/Q in coastal and open-ocean surface waters of the Northern Gulf of Mexico and U.S. East Coast off New Jersey. In situ measurements of scattering (b) from an ac9 probe were collected with concurrent measurements of bb from a six channel Hydrosat sensor. We compare the measured bb/b values with values derived from the Petzold Volume Scattering Functions (Petzold14), examine the spectral variability of the ratio and examine the variability in different water types related to the changes in the Volume Scattering Function (VSF). In addition, we estimate the T*f/Q (Mobley9) term from above-water measurements of remote sensing reflectance (Rrs) coupled with direct measurements of absorption (a) and backscattering (bb) coefficients. We will examine the spectral dependence of the T*f/Q term and its relationship to the bb/b ratio, which we use as a substitute for the changing VSF. Finally, we will show how the estimated T*f/Q values vary from the commonly used value of 0.051 used for satellite processing.					
15. SUBJECT TERMS backscattering, F/Q/ ratios, natural waters					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT SAR	18. NUMBER OF PAGES 9	19a. NAME OF RESPONSIBLE PERSON Bob Arnone
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified			19b. TELEPHONE NUMBER (Include area code) 228-688-5268

PUBLICATION OR PRESENTATION RELEASE REQUEST

Pubkey: 3366

NRLINST 5600.2

1. REFERENCES AND ENCLOSURES	2. TYPE OF PUBLICATION OR PRESENTATION	3. ADMINISTRATIVE INFORMATION
Ref: (a) NRL Instruction 5600.2 (b) NRL Instruction 5510.40D Encl: (1) Two copies of subject paper (or abstract)	<input type="checkbox"/> Abstract only, published <input type="checkbox"/> Book <input type="checkbox"/> Conference Proceedings (refereed) <input type="checkbox"/> Invited speaker <input type="checkbox"/> Journal article (refereed) <input type="checkbox"/> Oral Presentation, published <input type="checkbox"/> Other, explain <input type="checkbox"/> Abstract only, not published <input type="checkbox"/> Book chapter <input checked="" type="checkbox"/> Conference Proceedings (not refereed) <input type="checkbox"/> Multimedia report <input type="checkbox"/> Journal article (not refereed) <input type="checkbox"/> Oral Presentation, not published	STRN <u>NRL/PP/7330-02-62</u> Route Sheet No. <u>7330/</u> Job Order No. _____ Classification <u>X</u> U _____ C Sponsor _____ approval obtained <u>X</u> yes _____ no

4. AUTHOR

Title of Paper or Presentation

Variability in the Backscattering to Scattering and F/Q Ratios Observed in Natural Waters

Author(s) Name(s) (First, MI, Last), Code, Affiliation if not NRL

 Sherwin D. Ladner, Robert A Arnone, Richard W. Gould, Alan D. Weidemann, Vladimir I. Haltrin, Z. P. Lee, Paul M. Martinolich,
 Trisha Bergmann
It is intended to offer this paper to the Ocean Optics XVI

(Name of Conference)

18-NOV - 22-NOV-2002, Santa Fe, New Mexico, Unclassified

(Date, Place and Classification of Conference)

and/or for publication in Ocean Optics XVI, Unclassified

(Name and Classification of Publication)

(Name of Publisher)

After presentation or publication, pertinent publication/presentation data will be entered in the publications data base, in accordance with reference (a).

It is the opinion of the author that the subject paper (is _____) (is not X) classified, in accordance with reference (b).This paper does not violate any disclosure of trade secrets or suggestions of outside individuals or concerns which have been communicated to the Laboratory in confidence. This paper (does _____) (does not X) contain any militarily critical technology.This subject paper (has _____) (has never X) been incorporated in an official NRL Report.Robert A Arnone, 7333

Name and Code (Principal Author)

(Signature)

5. ROUTING/APPROVAL

CODE

SIGNATURE

DATE

COMMENTS

Author(s)

Haltrin

Section Head

Arnone

Branch Head

Steven W. Payne, 7330

Division Head

William J. Jobst, 7300

Security, Code

7030.1

Office of Counsel, Code

1008.3

ADOR/Director NCST

E.O. Hartwig, 7000

Public Affairs (Unclassified/
Unlimited Only), Code

7030.4

Division, Code

Author, Code

☒ A - Approved for public release, distribution is unlimited.

☐ B - Distribution authorized to U.S. Government agencies only (check reason below):

☐ Foreign Government Information

☐ Contractor Performance Evaluation

☐ Critical Technology

☐ Proprietary Information

☐ Administrative/Operational Use

☐ Premature Dissemination

☐ Test and Evaluation

☐ Software Documentation

☐ Cite "Specific Authority" _____
(Identification of valid documented authority)

Date statement applied _____

Other requests for this document shall be referred to _____
(Insert Controlling DOD Office*)

☐ C - Distribution authorized to U.S. Government agencies and their contractors (check reason below):

☐ Foreign Government Information

☐ Software Documentation

☐ Administrative/Operational Use

☐ Critical Technology

☐ Cite "Specific Authority" _____
(Identification of valid documented authority)

Date statement applied _____

Other requests for this document shall be referred to _____
(Insert Controlling DOD Office*)

☐ D - Distribution authorized to DOD and DOD contractors only (check reason below):

☐ Foreign Government Information

☐ Critical Technology

☐ Software Documentation

☐ Cite "Specific Authority" _____
(Identification of valid documented authority)

☐ Administrative/Operational Use

Date statement applied _____

Other requests for this document shall be referred to _____
(Insert Controlling DOD Office*)

☐ E - Distribution authorized to DOD components only (check reason below):

☐ Proprietary Information

☐ Premature Dissemination

☐ Critical Technology

☐ Foreign Government Information

☐ Software Documentation

☐ Direct Military Support

☐ Administrative/Operational Use

☐ Contractor Performance Evaluation

☐ Test and Evaluation

☐ Cite "Specific Authority" _____
(Identification of valid documented authority)

Date statement applied _____

Other requests for this document shall be referred to _____
(Insert Controlling DOD Office*)

☐ F - Further dissemination only as directed by _____
(Insert Controlling DOD Office*)

Date statement applied _____ or higher DOD authority _____

☐ G - Distribution authorized to U.S. Government agencies and private individuals or enterprises eligible to obtain export-controlled technical data in accordance with regulations implementing 10 U.S.C. 140c.

Date statement applied _____

Other requests for this document shall be referred to _____
(Insert Controlling DOD Office*)

*For NRL publications, this is usually the Commanding Officer, Naval Research Laboratory, Washington, DC 20375-5320

7. OTHER LIMITATION

☐ Classification

☐ NOFORN

☐ DTIC exempt (explain) _____

Classification Review
(initial/Date)

Substantive changes made in this document after approval by Classification Review and Public Release invalidate these reviews. Therefore, if any substantive changes are made by the author, Technical Information, or anyone else, the document must be returned for another Classification Review and Publication Release.

8. INSTRUCTIONS

Author completes and submits this form with the manuscript via line channels to the division head for review and approval according to the routing in Section 4.

1. NRL Reports.....Submit the diskette (if available), manuscript, typed double-spaced, complete with tables, illustrations, references, draft SF 298, and proposed distribution list.
2. NRL Memorandum Reports.....Submit a copy of the original, typed manuscript complete with tables, illustrations, references, draft SF 298, and proposed distribution list.
3. NRL Publications or other books, brochures, pamphlets,.....Handled on a per case basis by Site Technical Information Office. proceedings, or any other printed publications.

VARIABILITY IN THE BACKSCATTERING TO SCATTERING AND F/Q RATIOS OBSERVED IN NATURAL WATERS

Sherwin Ladner, Robert Arnone, Richard Gould*, Alan Weidermann*,
Vladimir I. Haltrin*, Zhongping Lee*, Paul Martinolich**, Trisha Bergmann****

*Planning Systems Incorporated, NRL Ocean Optics Section Code 7333,
Building 1009 RM 127A, Stennis Space Center, MS 39529, USA
Phone: (228) 688-5754, E-mail: ladner@nrlssc.navy.mil*

** Naval Research Laboratory, Ocean Optics Section, Code 7333,
Stennis Space Center, MS 39529, USA
E-mail: arnone@nrlssc.navy.mil*

*** Neptune Sciences Incorporated, 40201 Highway 190 East,
Slidell, LA 70461, USA
E-mail: martinol@nrlssc.navy.mil*

**** Rutgers University, Institute of Marine and Coastal Sciences,
New Brunswick, NJ 08901, USA
E-mail: bergmann@imcs.rutgers.edu*

ABSTRACT

Variability of the backscattering to scattering (b_b/b) and the upwelling irradiance/upwelling radiance (f/Q) ratios affect the accurate retrieval of inherent optical properties from ocean color satellite algorithms (SeaWiFS and MODIS). We investigate the variability of b_b/b and the f/Q in coastal and open-ocean surface waters of the Northern Gulf of Mexico and U.S. East Coast off New Jersey. In situ measurements of scattering (b) from an ac9 probe were collected with concurrent measurements of b_b from a six channel Hydrosat sensor. We compare the measured b_b/b values with values derived from the Petzold Volume Scattering Functions (Petzold¹⁴), examine the spectral variability of the ratio and examine the variability in different water types related to the changes in the Volume Scattering Function (VSF). In addition, we estimate the T^*f/Q (Mobley⁹) term from above-water measurements of remote sensing reflectance (R_{rs}) coupled with direct measurements of absorption (a) and backscattering (b_b) coefficients. We will examine the spectral dependence of the T^*f/Q term and its relationship to the b_b/b ratio, which we use as a substitute for the changing VSF. Finally, we will show how the estimated T^*f/Q values vary from the commonly used value of 0.051 used for satellite processing.

INTRODUCTION

Remote sensing algorithms for ocean color visible imagery are based on relationships between remote sensing reflectance and such inherent optical properties as absorption and scattering. Previous studies have simplified these relationships and are

currently used in present ocean color algorithms such as with SeaWiFS. The limitations of the relationships in different optical water types are under investigation. This effort is directed at examining different water types defined by the backscattering to scattering ratio and the changes in the T^*f/Q term (where Q is the ratio of upwelling irradiance to upwelling radiance). Both the ratios affect the accurate retrieval of the inherent optical properties.

The remote sensing reflectance (R_{rs}) is dependent on the angular scattering described by the volume scattering function (VSF). The VSF describes the angular distribution of scattered light in the water column. The integral of the VSF over full solid angle is the scattering coefficient (b); the integral over angles in the backward direction only (90-180 degrees) is the backscattering coefficient (b_b). The VSF is influenced by particle characteristics such as shape, index of refraction, and size distribution. Because the VSF is difficult to measure (new instruments are just now becoming available), we use the b_b/b ratio (or probability of backscattering) as a substitute for the VSF to describe water mass differences. This ratio provides a method to describe the differences in the VSF for different water masses and how the angular differences influence the R_{rs} . Understanding the differences in b_b/b for different water masses will help characterize the changes in the VSF, which affect R_{rs} estimates, and provide better estimates of inherent optical properties in coastal and open-ocean waters (Northern Gulf of Mexico and New Jersey Coast) where different b_b/b water types are observed. This is important for researchers to model optical remote sensing algorithms, visibility and laser propagation in seawater.

We will use in situ measurements for various water types to determine how the b_b/b changes both spectrally and in a wide variety of coastal and open-ocean waters and how it compares with the linear Petzold¹⁴ b_b/b ratio. Our second objective is to evaluate the relationship between measured remote sensing reflectance, backscattering and absorption through the ratio $[b_b/(a+b_b)]$. We will examine the spectral nature of the T^*f/Q term and show how the estimated T^*f/Q values vary from the commonly used value of 0.051 used for satellite processing

BACKGROUND

The availability of VSF, b_b and b measurements has been limited in the past. VSF measurements (515nm) made in San Diego Harbor over 30 years ago by Petzold¹⁴ were used by Gould et al.⁵ to estimate a linear relationship between b_b and b :

$$b_b = 0.01829 * b + 0.00006, \quad (r^2 = 0.99) \quad (1)$$

This is a limited data set of 15 angular scattering coefficients and attenuation coefficients of 15 phase functions, collected at a single wavelength that does not describe the differences in the VSF for different water types. New instruments are currently available that measure the VSF (Haltrin et al.⁶), backscattering and scattering directly.

Irradiance reflectance or diffuse reflectance coefficient is defined as:

$$R = E_u/E_d \quad (2)$$

Using different approaches, Morel and Prieur¹³ and Gordon et al.² related irradiance reflectance (R) to water inherent optical properties (absorption and backscattering) through a factor f :

$$R = f*[b_b/(a+b_b)] \quad (3)$$

where f varies in a range of 0.28 – 0.42 for most “Case 1” waters and for solar zenith angles less than 70 degrees (Morel and Gentili¹⁰). Remote sensing reflectance is defined as:

$$R_{rs} = L_w/E_d \quad (4)$$

To convert R to R_{rs} (in order to relate satellite measurements to in-water optical properties), we must apply an irradiance-to-radiance conversion factor Q (L_w/E_d) and push the radiance/irradiance values through the air/sea interface. Following notation in Mobley⁹:

$$R_{rs} = (T/Q) * R \quad (5)$$

(See Mobley⁹ for derivation of T term). Substituting for R from equation 3

$$R_{rs} = (T*f/Q) * [b_b/(a+b_b)] \quad (6)$$

The value of T is approximately 0.54 and only varies by a few percent (Mobley⁹). From a theoretical perspective, f/Q varies with sun and satellite viewing angle and changing inherent optical properties and ranges from 0.07 to 0.11 over realistic angles for remote sensing applications (Morel and Gentili¹¹). It changes in different water types in response to VSF and angular scattering. So considering the variability in both T and f/Q , we would expect a range of approximately 0.0366 to 0.0612 for the $T*f/Q$ term in equation 6. Using in situ measurements, we describe how this term changes spectrally and in different water masses.

Lee et al.⁷ are further refining the estimation of f/Q by analyzing and modeling the effect of the VSF shape on this term. The new model, weighted by relative contribution of particle scattering, provides improved estimations of f/Q . This refined estimation of f/Q will further improve estimations of R_{rs} and in-water properties, but does require a priori knowledge of a and b_b .

METHODS

We collected data at 174 stations in different waters types (open ocean and coastal). These included two cruises in the Northern Gulf of Mexico in May 2001 and May 2002 off of Mobile Bay for a total of 83 stations. The remaining 91 stations were collected in New Jersey in July 2000 and July 2001. The Gulf of Mexico represents a wide variety of highly scattering waters with elevated suspended sediment loads and high CDOM where as the New Jersey cruises represent clearer waters with less sediments. We collected above-water measurements of R_{rs} using an Analytical Spectral Device

(ASD) field spectroradiometer which was processed using near-infrared (NIR) surface glint removal algorithms (Gould et al.⁵). This is required to account for the Rrs at near-infrared wavelengths in high scattering waters. The ASD measures the spectrum at 1.3 nm resolution from 400 – 800 nm and a 12% spectralon (gray) card was used for calibration. Absorption and beam attenuation (a and c) were measured using a WetLabs ac9 at 9 wavelengths (412, 440, 488, 510, 532, 555, 650, 676, 715nm). The instrument was calibrated using milli-Q water and the Zaneveld scatter correction was applied (Zaneveld et al.¹⁶). We derive scattering by difference ($c-a$). The spectral backscattering coefficient was measured with the Hydroscat instrument, which measures the scattering at 140 degrees and extrapolates the b_b at 6 wavelengths (442, 488, 532, 589, 620, 676nm) (Maffione and Dana⁸). The instrument was calibrated at the factory prior to the deployments. We removed the backscattering and scattering due to pure water using Smith and Baker¹⁵ to investigate the differences in the spectral b_b/b ratio.

RESULTS

Backscattering to Scattering Relationship

We plotted measured b_b and b for two regions (Gulf of Mexico and New Jersey) to investigate the relationship for 440, 488, 532, and 676nm and differences for different water types. Figure 1 shows the relationship between b_b and b at 532nm for the Gulf of Mexico 2001 and 2002 including Petzold¹¹ data at 515nm. The Gulf of Mexico data seems to depict two different water types, although there is much spread in the data and considerable overlap. The first water type includes Gulf of Mexico data from the 2002 cruise (triangles and dotted linear fit) and is similar to the Petzold¹⁴ relationship (open circles). The second water type is defined by the Northern Gulf of Mexico from 2001 cruise (filled circles and solid linear fit). Both Northern Gulf of Mexico cruises were conducted in the same vicinity one year apart.

Figure 2 shows the relationship between b_b and b at 532nm for the New Jersey cruises. The data displayed in Figure 2 more clearly depict two distinct water types. The first water type includes data from the 2000 cruise (filled circles and solid linear fit) and is similar to the Petzold relationship (open circles). The second water type is defined by the data from the 2001 cruise (triangles and dotted linear fit). This water type consist of mostly open ocean waters dominated by biological rather than inorganic particles and produced a b_b/b ratio lower than the other cruises. Note that the New Jersey cruises were also conducted in same vicinity one year apart.




Figure 1. Measured b_b (532nm) vs. b (532nm) for the two Gulf of Mexico Cruises




Figure 2. Measured b_b (532nm) vs. b for the two New Jersey Cruises.

By using the slopes of the lines in figures 1 and 2, which is essentially the b_b/b ratio for all stations in the relationship, and using the other measured wavelengths (440,488,676) in common between the two instruments, we can examine the spectral dependence of the b_b/b ratio. Figures 3 and 4 represent the b_b/b ratios derived from Figures 1 and 2 plotted as function of wavelength. Figure 3 shows the two distinct water types for the Northern Gulf of Mexico region. Figure 4 shows the two distinct water types for the New Jersey region. The spectral shape of both regions is nearly flat. We will further examine the statistical difference at a later time. The Northern Gulf of Mexico waters during 2002 and the New Jersey 2000 waters are similar to that of Petzold¹⁴ type waters. We note that for the New Jersey waters the spectral shapes are somewhat similar and the water types show more distinct difference than the Northern Gulf of Mexico cruises. Results show the b_b/b relationship is not always related to the Petzold¹⁴ relationship. The changes in the b_b/b ratio can possibly be used to characterize the VSF in different water types and agrees with the behavior seen in actual VSF measurements (Haltrin et al.⁶).




Figure 3. Measured b_b/b ratio (y-axis) as a function of wavelength (x-axis) for two Gulf of Mexico Cruises.




Figure 4. Measured b_b/b ratio (y-axis) as a function of wavelength (x-axis) for two New Jersey Cruises.

F/Q Ratio

We took the measured properties of a , b_b , and computed the ratio $[b_b/(a+b_b)]$ for the Gulf of Mexico and New Jersey stations combined (Figure 5). The regions were plotted separately (not shown) but did not show a regional dependence for this study. We plotted the relationship between the measured R_{rs} at 440 (top left), 488 (top right), 532 (bottom left) and 676nm (bottom right) versus the $[b_b/(a+b_b)]$ ratio. These relationships show the spectral variance in the (T^*f/Q) term used in equation 6 for these two regions. From theoretical studies, this term should range between approximately 0.0366 – 0.0612. Based on measured values, our results show that the term ranged from 0.04 (532nm) - 0.07 (440nm). A value of 0.051 is commonly used in satellite ocean color processing (Carder et al.¹). The 532nm (lower left) channel shows higher scatter than the others. This scatter could be because the 532nm channel has a higher in-water transparency than other channels and is where the minimum absorption occurs. Any small variation in absorption, whether it is due to the error in measurement or stability of the instrument, could possibly cause such high variations for this relationship.




Figure 5. Measured R_{rs} vs. $b_b/(a+b_b)$ ratio for all stations collected in the Northern Gulf of Mexico and New Jersey regions.

We took the mean of the T^*f/Q terms estimated by dividing measured reflectance by the measured ratio $[b_b/(a+b_b)]$ and plotted as a function of wavelength shown in figure 6. Morel and Gentili¹¹ also looked at the f/Q term as a function of wavelength. They found that the values for f/Q to be 0.0936 (440nm), 0.0944 (500nm), 0.0929 (565nm), and 0.0881 (665nm), which would result in T^*f/Q estimates of 0.0505, 0.0509, 0.0502, and 0.0476. Their results are stable and linear as a function of wavelength whereas results in this study show a larger variation. Morel and Gentili¹¹ observed Case I waters and radiances encountered in the vicinity of the vertical direction. The measurements of R_{rs} in this study were measured 40 degrees from nadir and collected in non-Case I waters. Morel and Gentili¹¹ documented that before interpreting the marine signals, the discrimination between Case I and sediment-dominated Case II waters has to be made, therefore a separate knowledge of the Q factor is needed.




Figure 6. Mean T^*f/Q calculated from equation 6 vs. wavelength. Error Bars indicate + or - one standard deviation.

CONCLUSION

The b_b/b ratio derived from in situ measurements varies spectrally over different water types (mostly turbid). This variability maybe related to regional differences in the VSF. We looked at four different cruises (two in the Northern Gulf of Mexico and two off the coast of New Jersey). We clearly observed two different water types for each region but the b_b/b ratio as a function of wavelength was nearly flat spectrally. In some cases for the Gulf of Mexico and New Jersey region the b_b/b ratios were quite similar to the ones derived from the Petzold¹⁴ data while in others it was not. This ratio provides a method to describe the differences in the VSF for different water masses and how the angular differences influence the R_{rs} .

We estimated the T^*f/Q term from above-water measurements of R_{rs} coupled with direct measurements of absorption and backscattering. We show in this study that the estimated T^*f/Q term in non-Case I waters has more spectral variability compared to

values presented by Morel and Gentili¹¹ in Case I waters. From theoretical considerations, T^*f/Q from equation 6 should range between 0.0366 and 0.0612. In our current satellite processing we use a constant value for T^*f/Q equal to 0.051. Based on in situ measurements, we found the spectral range of the term to be between 0.04 (532nm) and 0.086 (440nm) for the four wavelengths examined in this study. We did not note any separation due to water type when looking at the four cruises.

The understanding of the differences in the b_b/b relationship will help characterize the changes in the VSF, which affect R_{rs} estimates, and the parameterization of the f/Q term for different water types certainly can improve the accuracy of algorithms applied to remotely sensed ocean color data.


ACKNOWLEDGEMENTS

Support for this work was provided by the Naval Research Laboratory through (Program Element 0601153N). This article represents NRL contribution NRL/PP/7330-02-62.

REFERENCES

1. K. L. Carder, F. R. Chen, Z. P. Lee, S. K. Hawes, and K. Kamykowski, Semianalytic Moderate-Resolution Imaging Spectrometer algorithms for chlorophyll a and absorption with bio-optical domains based on nitrate-depletion temperatures, *JGR*, 104, C3, 5403-5421, 1999.
2. H. R. Gordon, O. B. Brown, R. H. Evans, J. W. Brown, R. C. Smith, K. S. Baker, and D. K. Clark, A semi-analytical radiance model of ocean color, *JGR*, 93, 10909-10924, 1988
3. H. R. Gordon, O. B. Brown, and M. M. Jacobs, Computed relationships between the inherent and apparent optical properties of a flat homogeneous ocean, *Appl. Opt.*, 14, 417-427, 1975.
4. R. W. Gould, R. A. Arnone, and P. M. Martinolich, Spectral dependence of the scattering coefficient in case 1 and case 2 waters, *Appl. Opt.* 38, 12, 2377-2383, 1999.
5. R. W. Gould, R. A. Arnone, and M. Sydor, Absorption, scattering and remote-sensing reflectance relationships in coastal waters: Testing an new inversion algorithm, *Journal of Coastal Research*, 17, 2, 328-341, 2001.
6. V. I. Haltrin, M. E. Lee, E. B. Shybanov, R. A. Arnone, A. D. Weidemann, V. I. Mankovsky, W. S. Pegau, and S. D. Ladner, Relationship between backscattering and beam scattering coefficients derived from new measurements of light scattering phase functions, *Ocean Optics XVI Proceedings*, 2002.
7. Z. P. Lee, K. L. Carder, and K. P. Du, Influence of particle phase functions on the the model parameter of remote-sensing reflectance, *Ocean Optics XVI Proceedings*, 2002.

8. R. A. Maffione and D. R. Dana, Instruments and methods for measuring the backward-scattering coefficient of ocean waters, *Appl. Opt.* 36, 24, 6057-6067, 1997.
9. C. D. Mobley, Estimation of the remote-sensing reflectance from above-surface measurements, *Appl. Opt.* 38, 36, 7442-7454, 1999.
10. A. Morel and B. Gentili, Diffuse reflectance of oceanic waters: its dependence on Sun angles as influenced by the molecular scattering contribution, *Appl. Opt.* 30, 4427-4438, 1991.
11. A. Morel and B. Gentili, Diffuse reflectance of oceanic waters, II, Bidirectional Aspects, *Appl. Opt.*, 32, 6864-6879, 1993.
12. A. Morel and B. Gentili, Diffuse reflectance of oceanic waters, III, Implication of bidirectionality for the remote-sensing problem, *Appl. Opt.* 35, 24, 4850-4862, 1996.
13. A. Morel and L. Prieur, Analysis of variations in ocean color, *Limnol. Oceanogr.*, 22, 709-722, 1977.
14. T. J. Petzold, Volume scattering functions for selected ocean waters, *Scripps Institute of Oceanography Ref.* 72-78, 79, 1972
15. R. C. Smith and K. S. Baker, Optical properties of the clearest natural waters (200-800nm), *Appl. Opt.* 20, 177-184, 1981.
16. J. R. V. Zaneveld, J. C. Kitchen, and C. Moore, The scattering error correction of reflecting-tube absorption meters, *Ocean Opt. XII Proc.*, SPIE 2258, 44-55, 1994.



Program and Abstracts

COVERING A DIVERSITY OF TOPICS PERTAINING TO:

- Radiative Transfer Theory
- In Situ and Remote Sensor and Measurement System Design
 - Applications of Quantitative Remote Sensing
- Interdisciplinary Processes Pertaining to the Ocean Surface, Volume, and Floor
 - Marine Environment Management Strategies

Sponsored by:

Office of Naval Research

United States Navy and Space Administration